

Research Article

Maternal Risk Factors Associated with Neonatal Stunting: A Case–Control Study

Anita Rohmah¹, Yasmini Fitriyati², Afifah Az Zahra³, Isna Arifah Rahmawati³,
Muhtia Mahderina Cahyani³

¹Department of Obstetrics and Gynecology Wonosari Hospital

²Department of Obstetrics and Gynecology

³Faculty of Medicine, Universitas Islam Indonesia
Yogyakarta

Abstract

Objective: To identify maternal risk factors associated with neonatal stunting at Wonosari Hospital.

Methods: A case–control study was conducted involving mothers who gave birth at Wonosari Hospital in 2023. Maternal sociodemographic characteristics (age, education, and occupation), nutritional status (body mass index and mid-upper arm circumference), and pregnancy-related factors (gestational age, gestational status, hemoglobin levels, blood pressure, pregnancy complications, and mode of delivery), as well as newborn length at birth, were obtained from medical records. Neonatal stunting was defined as a length-for-age z-score < -2 SD according to World Health Organization (WHO) standards. Statistical analysis was performed using chi-square tests and multiple logistic regression.

Results: A total of 154 participants were included, equally divided into case (stunted newborns) and control (non-stunted newborns) groups. Mothers with a lower educational level had significantly higher odds of delivering a stunted newborn ($p = 0.010$; aOR = 2.845; 95% CI = 1.286–6.293). Preterm birth was also associated with an increased risk of neonatal stunting ($p = 0.033$; aOR = 9.847; 95% CI = 1.210–80.152). In addition, pregnancy complications were significantly associated with higher odds of neonatal stunting ($p = 0.020$; aOR = 2.728; 95% CI = 1.171–6.352).

Conclusion: Maternal factors, including low educational level, preterm birth, and pregnancy complications, were significantly associated with neonatal stunting at Wonosari Hospital. These findings underscore the importance of maternal education in neonatal health outcomes. Furthermore, close monitoring of fetal growth and nutritional status, along with appropriate management of pregnancy complications, may help reduce the risk of neonatal stunting. However, larger-scale studies are needed to assess the population-level impact of these factors.

Keywords: neonatal, pregnancy complication, stunting.

Correspondence author. Afifah Az Zahra. Universitas Islam Indonesia. Yogyakarta,
Email: 217110403@uui.ac.id Telp : +6285729222530

INTRODUCTION

During the first 1000 days of life, the period from conception to the first two years of a child's life, maternal nutrition plays a crucial role in determining fetal health and future well-being. During pregnancy, the fetus undergoes rapid development and requires adequate nutrient intake from the mother. The increased nutritional needs of pregnant women are influenced by age, pre-pregnancy nutritional status, and any underlying medical conditions¹.

Pregnant women with malnutrition,

undernutrition, or poor nutritional status during pregnancy are more likely to give birth to infants with low birth weight or small for gestational age². Basic Health Research (Rskesdas), 48.9% of pregnant women were anemic, 17.3% experienced Chronic Energy Deficiency (CED), and 28% had a risk of life-threatening obstetric complications. Several studies have shown that deficiencies in specific nutrients during pregnancy can negatively impact fetal growth and contribute to low birth weight³. For high-risk pregnant women, preventive measures and nutritional interventions become increasingly

important to prevent complications and fetal developmental problems⁴.

Neonatal stunting is a growth disorder that characterized by a birth length-for-age z-score of <-2 Standard Deviations (SD)⁵. This condition is a serious global public health issue, particularly in developing countries like Indonesia. Global Nutrition Report, the prevalence of neonatal stunting reached 21.9% in low- and middle-income countries⁶. Stunting has long-term impacts on health, cognitive development, and quality of life, even into adulthood⁷.

Malnutrition in Indonesia remains one of the highest in the world, with 1 in 10 children under five experiencing wasting and 3 in 10 children experiencing stunting (short stature). Based on the results of the Indonesian Nutrition Status Survey (SSGI), the prevalence of stunting in Indonesia decreased by 2.8%, from 24.4% in 2021 to 21.6% in 2022. This achievement is in line with the target set by the Ministry of Health, which is approximately 2.7% annually. This study aims to examine the characteristics of pregnant women associated with neonatal stunting at a referral hospital in the Special Region of Yogyakarta, Wonosari Hospital.

METHODS

This observational analytical study employed a case-control design to investigate maternal characteristics associated with neonatal stunting. The study population consisted of pregnant women who delivered at Wonosari Hospital among March and September 2023.

The sample size was calculated using the Lemeshow formula, with subjects distributed into two groups at a 1:1 ratio. The case group comprised mothers who delivered stunted newborns, while the control group comprised mothers who delivered newborns with normal birth length. Subjects were selected using purposive sampling to minimize selection bias. To control for information bias inherent in secondary data usage, strict inclusion and exclusion criteria were applied. Only medical records with complete documentation were included. Records with missing data on key variables, were excluded from the study.

The dependent variable was the incidence of neonatal stunting. In accordance with WHO growth standards and current clinical references, neonatal stunting was operationally defined as a birth length-for-age z-score of <-2 SD relative

to the WHO child growth standards. Data were collected retrospectively using secondary data from patient medical records. Data validity and reliability were ensured, as all anthropometric measurements (birth length, maternal Mid-Upper Arm Circumference/MUAC) and vital signs (blood pressure) were performed by certified healthcare professionals (midwives and nurses) using calibrated, hospital-standard instruments (infantometers for newborns and digital sphygmomanometers for mothers).

Independent variables encompassed maternal characteristics and health status, specifically: maternal age, educational attainment, employment status, parity, gestational age (preterm <37 weeks vs term ≥ 37 weeks), nutritional status (BMI and MUAC <23.5 cm for Chronic Energy Deficiency/CED), hemoglobin levels (anemia defined as <11 g/dL), blood pressure (hypertension defined as $\geq 140/90$ mmHg), and other concurrent pregnancy complications.

Statistical analysis was performed using IBM SPSS Statistics software version 25. Univariate analysis was utilized to describe the demographic characteristics of the subjects. Associations between variables were assessed using bivariate analysis (chi-square test). Variables meeting the inclusion criteria ($p < 0.25$) were subsequently entered into a multivariate analysis using multiple logistic regression to identify dominant risk factors. This study received ethical clearance from the Medical Research Ethics Committee of the Faculty of Medicine, Universitas Islam Indonesia with the registered number: 18/Ka.Kom.Et/70/KE/XII/2023).

RESULTS

This study enrolled 154 participants, equally distributed into case and control groups ($n=77$ each). The majority of subjects (Table 1) fell within the healthy reproductive age range of 20–34 years ($n=121$, 78.6%) and exhibited normal nutritional status ($n=117$, 76%). High prevalence of pregnancy complications (74.7%) reflects the study setting at Wonosari Hospital as a referral center, where cases were predominantly characterized by severe preeclampsia ($n=50$, 32.5%) and premature rupture of membranes (PROM) ($n=27$, 17.5%), alongside other conditions such as oligohydramnios and gestational diabetes.

Table 1. Baseline Characteristics of Mothers at Wonosari Hospital

Characteristic	N	(%)
Age (y o)		
<20	5	3.2
20 – 35	121	78.6
≥35	28	18.2
Education		
Elementary	46	29.9
Secondary	104	67.5
High	4	2.6
Working Status		
Yes	50	32.5
No	104	67.5
Nutritional Status		
Underweight	5	3.2
Normal	117	76
Overweight	27	17.5
Obesity	5	3.2
History of PEM		
Yes	11	7.1
No	143	92.9
Anemia		
Yes	30	19.5
No	124	80.5
Gestational Hypertension		
Yes	72	46.8
No	82	53.2
Gestational Age		
Preterm	13	8.4
Aterm	141	91.6
Number of Pregnancies		
Primigravida	56	36.4
Multigravida	98	63.6
Complications		
Yes	115	74.7
No	39	25.3
Childbirth		
Normal	81	52.6
C-section	73	47.4

The association between maternal characteristics and neonatal stunting was assessed using the chi-square test (Table 2). Preterm birth (<37 weeks) significantly increased the risk of neonatal stunting ($p=0.012$; $OR=14.031$; 95%CI:1.776–110.824). Mothers experiencing pregnancy complications had a fourfold greater risk of delivering a stunted newborn compared to those without complications ($p=0.001$; $OR=4.048$; 95%CI:1.803–9.087). Additionally, mothers with a basic education level demonstrated a significantly higher proportion of stunted newborns compared to those with a secondary education ($p=0.001$).

Other variables, including maternal age, employment status, nutritional status (BMI), history of Chronic Energy Deficiency (CED), anemia, parity, and history of hypertension, did not show statistically significant associations with neonatal stunting. Although maternal anemia appeared to nearly double the risk of stunting, statistical analysis indicated that this association was not significant ($p=0.107$; $OR=1.966$; 95%CI:0.864–4.472).

Table 2. Bivariate Analysis of Maternal Risk Factors Associated with Neonatal Stunting

Characteristic	Neonatal		OR (CI 95%)	P-value
	Stunted	Normal		
Age (y o)				
<25 and >35	20	60.6	13	39.4
20-34	57	47.1	64	52.9
			0.935 (0.454-1.922)	0.851
Education				
Elementary	20	60.6	16	39.4
Secondary	57	47.1	64	52.9
High	1	25	3	75
			Ref 3.601 (1.70-7.63) 7.615 (0.73-80.05)	0.001 0.091
Working status				
Yes	22	44	28	56
No	55	53	49	47
			0.700 (0.355-2.379)	0.302
Nutritional status				
Underweight	4	80	1	20
			0.238 (0.26-2.189)	0.205

Normal	57	48.7	60	51.3	Reff	
Overweight	14	52	13	48	0.882 (0.382-2.038)	0.769
Obesity	2	40	3	60	1.425 (0.230-8.844)	0.704
History of PEM						
Yes	6	54.5	5	45.5	1.217 (0.355-4.169)	0.755
No	71	49.7	72	50.3		
Anemia						
Yes	19	63.3	11	36.7	1.966 (0.864-4.472)	0.107
No	58	46.8	66	53.2		
Gestational Hypertension						
Yes	37	51.4	35	48.6	0.901(0.478-1.697)	0.747
No	40	48.8	42	51.2		
Gestational Age						
Preterm	12	92.3	1	7.7	14.031 (1.776-110.824)	0.012
Aterm	65	46.1	76	53.9		
Number of pregnancy						
Primigravida	27	48.2	29	51.8	0.894 (0.463-1.724)	0.738
Multigravida	50	51	48	49		
Complication						
Yes	67	58.3	48	41.7	4.048 (1.803-9.087)	0.001
No	10	25.6	29	74.4		

Multiple logistic regression analysis was performed on maternal risk factors including gestational age, pregnancy complications, maternal education level, and anemia ($p<0.25$). As detailed in Table 3, the final model identified three primary predictors. Gestational Age emerged as the strongest risk factor, where preterm birth increased the risk of neonatal stunting by 9.8

times ($p=0.033$; aOR=9.847; 95%CI:1.210–80.152). Maternal Education was a significant predictor ($p=0.010$), with basic education associated with a higher risk compared to secondary education (aOR=2.845; 95%CI:1.286–6.293). Pregnancy Complications independently increased the risk of neonatal stunting by 2.7 times ($p=0.020$; aOR=2.728; 95%CI:1.171–6.352).

Table 2. Bivariate Analysis of Maternal Risk Factors Associated with Neonatal Stunting

Variabel	B	S.E	Wald	df	aOR (aCI 95%)	P-value
Education Level			7.422	2		0.024
Secondary	1.046	0.405	6.666	1	2.845 (1.286-6.293)	0.010
High	1.767	1.218	2.103	1	5.851 (0.537-63.698)	0.147
Gestational Age	2.287	1.070	4.571	1	9.847 (1.210-80.152)	0.033
Maternal Complication	1.003	0.431	5.413	1	2.728 (1.171-6.352)	0.020
Constant	-6.449	2.146	9.029	1	0.002	0.003

The analysis confirms that all three variables gestational age, maternal education, and pregnancy complications are significant independent risk factors for neonatal stunting. This is evidenced by adjusted odds ratio (aOR) values greater than 1 for each variable. Gestational age shows the highest point estimate (aOR = 9.85), although its wide confidence interval reflects considerable uncertainty, likely due to the limited number of preterm cases. In contrast, maternal education and pregnancy complications present narrower confidence intervals, indicating more precise and stable risk estimates within the studied population.

DISCUSSION

This study confirms that maternal education is a critical determinant of neonatal health. We found that lower maternal education is significantly associated with an increased risk of delivering stunted infants ($p=0.010$; aOR=2.845; 95%CI=1.286–6.293). This results is consistent with a study on stunting determinants which highlighted that limited health literacy and delayed healthcare seeking among mother with lower education directly compromise maternal nutrient intake⁸. Higher maternal education and literacy correlate with improved socioeconomic status and household income, thereby facilitating access to essential resources such as nutritious

food, healthcare, and sanitation while enhancing the understanding of child care practices required for optimal growth.^{9,10}

Therefore, preventing neonatal stunting requires more than clinical intervention; it necessitates targeted educational empowerment for mothers with lower formal education, specifically regarding the first 1000 days of life. This approach is supported by recent findings demonstrating that nutritional education significantly enhances maternal understanding of adequate nutrition from preconception through pregnancy, a factor proven to reduce stunting risks¹¹.

In this study, the majority of stunted newborns in this study were born prematurely (<37 weeks), establishing lower gestational age as a significant predictor of neonatal stunting ($p=0.03$; aOR 9.847; 95%CI 1.210-80.152). This finding is consistent with previous observations demonstrating that premature newborns tend to have lower birth weight and length compared to full-term counterparts¹². Growth impairment initiating at birth often persist through the first two years of life, predisposing the child to permanent short stature and a heightened risk of chronic comorbidities in adulthood¹³.

Underlying this growth restriction is the fact that bone mineralization is critically dependent on the third trimester of gestation, during which approximately 80% of fetal mineral accretion primarily calcium and phosphate occurs via active placental transport¹⁴. Consequently, preterm birth abruptly interrupts this vital supply, leaving the infant with significant mineral deficits. Furthermore, at the molecular level, this growth restriction is exacerbated by disruptions in the Growth Hormone (GH) axis, characterized by suppressed Insulin-like Growth Factor 1 (IGF-1) concentrations¹⁵. Systemic inflammation significantly worsens this deficiency, as proinflammatory cytokines drive the upregulation of IGFBP-1, which sequesters free IGF-1 and limits its bioavailability for linear bone mineralization¹⁶. Concurrently, nutritional deficits and intrauterine growth restriction are linked to elevated IGFBP-2 levels, providing a distinct pathway of growth inhibition that compounds the effects of inflammation¹⁷.

This study demonstrates that pregnancy complications, particularly severe preeclampsia and premature rupture of membranes (PROM), significantly increase the risk of neonatal stunting ($p=0.020$; aOR=2.728; CI=1.171-6.352).

The mechanism linking these complications to stunting likely involves placental insufficiency and inflammation. Preeclampsia and gestational hypertension induce systemic vasoconstriction and placental hypoperfusion, restricting the transfer of oxygen and nutrients essential for fetal skeletal growth¹⁸⁻²⁰.

Similarly, PROM is often precipitated by ascending urinary tract infections or systemic inflammation, which can trigger preterm labor and interrupt the fetal development timeline^{21,22}. While other studies have linked factors sleep disturbances²³, HIV²⁴, and gestational diabetes^{25,26} to adverse outcomes, our findings specifically underscore that in this population, aggressive management of hypertension and infection control is the direct preventive strategy for neonatal stunting. Consequently, stratified antenatal care focusing on these specific pathological drivers is essential to mitigate the risk of neonatal growth failure.

Several sociodemographic variables traditionally cited as risk factors did not demonstrate statistical significance in this study. In contrast to previous findings linking reproductive extremes (<20 or >35 years) to stunting¹², our analysis revealed no such association. While these age groups are recognized as high-risk for pregnancy complications^{27,28}, our results align with other studies suggesting that stunting is driven more by modifiable determinants such as maternal education and access to prenatal care rather than biological age alone²⁹.

Similarly, the lack of association with gravidity in our study is consistent with research suggesting that maternal nutritional reserves and socioeconomic status are more critical predictors than the number of pregnancies³⁰, although short inter-pregnancy intervals (<24 months) remain a potential risk factor³¹. Furthermore, employment status was not a significant differentiator. While excessive workload (>40-hour shifts) is a known stressor^{32,33}, occupation itself is likely a secondary factor that can be modified to reduce morbidity rather than a primary cause of stunting³⁴.

Maternal nutritional status (BMI and MUAC) was not significantly associated with neonatal stunting, a finding that contradicts established evidence linking maternal underweight status to fetal growth restriction and micronutrient deficiencies (iron, folate, zinc)^{35,36}. Previous studies emphasize that maternal height (<150 cm) and weight (<43 kg) are strong predictors of stunting³⁷. Nutrition impairment

such as malnutrition hinders fetal growth through nutrient deprivation^{36,38}, while obesity impairs placental perfusion with inflammatory pathways^{19,39}. The lack of significance in our data may be attributed to sample limitations or the influence of unmeasured confounders often linked to low socioeconomic status, such as poor sanitation and low per capita income^{40,41} or behavioral factors like smoking, alcohol, caffeine intake and substance exposure^{42,43}.

Maternal anemia did not reach statistical significance ($p=0.107$). However the analysis revealed a clinically relevant trend, with anemic mothers having nearly twice the risk of delivering stunted newborns ($OR=1.966$). This contradicts findings where anemia showed a stronger statistical impact on birth length^{44,45}. Anemia induces chronic fetal hypoxia and restricts the oxygen supply essential for development, often resulting in congenital malformations or growth restriction^{46,47}. However, the absence of statistical significance in our dataset aligns with recent studies in developing countries, which suggest that neonatal stunting may be more strongly correlated with maternal short stature and chronic deprivation rather than acute anemia levels during pregnancy alone^{12,48}.

This study is limited by its retrospective design and reliance on secondary data, which may introduce selection bias and reduce the accuracy of some measurements. The single-hospital setting restricts generalizability, while the small sample size limited statistical power, particularly in the underweight subgroup. Finally, unmeasured confounders like paternal height and daily nutritional intake were not included in the analysis.

CONCLUSION

This study concludes that low maternal education, preterm birth, and pregnancy complications significantly increase the risk of neonatal stunting. However, maternal nutritional status and anemia did not demonstrate statistical significance in this study. Strengthening maternal education and improving antenatal care services are essential in reducing stunting risk, particularly through stratified monitoring of maternal comorbidities and targeted health literacy programs. Further longitudinal studies are required to explore long-term child outcomes and causal mechanisms to inform public health policies.

ACKNOWLEDGMENT

The authors thank the Director and the Medical Records Department of Wonosari Hospital, Yogyakarta, for facilitating the data collection process. We also acknowledge the Faculty of Medicine, Islamic University of Indonesia, for providing institutional support.

REFERENCES

1. Singh DP, Biradar RA, Halli SS, Dwivedi LK. Effect of maternal nutritional status on children nutritional status in India. *Child Youth Serv Rev.* 2021;120:105727. doi:10.1016/j.childyouth.2020.105727
2. Setiawan AS, Indriyanti R, Suryanti N, Rahayuwati L, Juniarti N. Neonatal stunting and early childhood caries: A mini-review. *Front Pediatr.* 2022;10. doi:10.3389/fped.2022.871862
3. Lassi ZS, Kedzior SGE, Tariq W, Jadoon Y, Das JK, Bhutta ZA. Effects of preconception care and periconception interventions on maternal nutritional status and birth outcomes in low-and middle-income countries: A systematic review. *Campbell Systematic Reviews.* 2021;17(2). doi:10.1002/cl2.1156
4. Saleh A, Syahrul S, Hadju V, Andriani I, Restika I. Role of Maternal in Preventing Stunting: a Systematic Review. *Gac Sanit.* 2021;35:S576-S582. doi:10.1016/j.gaceta.2021.10.087
5. Kerac M, Ashorn P, Berkley JA, et al. Infants less than 6 months of age at risk of poor growth and development: evidence gaps identified during WHO guideline development. *BMJ Glob Health.* 2025;10(Suppl 5):e017227. doi:10.1136/bmjgh-2024-017227
6. Mason JB, Shrimpton R, Saldanha LS, et al. The first 500 days of life: policies to support maternal nutrition. *Glob Health Action.* 2014;7(1):23623. doi:10.3402/gha.v7.23623
7. Enlow MB, Sideridis G, Bollati V, Hoxha M, Hacker MR, Wright RJ. Maternal cortisol output in pregnancy and newborn telomere length: Evidence for sex-specific effects. *Psychoneuroendocrinol.* 2019;102:225-35. doi:10.1016/j.psyneuen.2018.12.222
8. Beal T, Tumilowicz A, Sutrisna A, Izwardy D, Neufeld LM. A review of child stunting determinants in Indonesia. *Matern Child Nutr.* 2018;14(4):e12617. doi:10.1111/mcn.12617
9. Munanadia M. Perilaku Ibu Hamil dalam Pencegahan Stunting di Puskesmas Panarung. *Bunda Edu-Midwifery J (BEMJ).* 2022;5(2):31-6. doi:10.54100/bemj.v5i2.66
10. Meshram II, Kodavanti MR, Krishna KS, Laxmaiah A. Regional Variation in the Prevalence of Undernutrition and its Correlates Among Under Five-Year Children in North India. *Indian J Comm Med.* 2024;49(2):322-33. doi:10.4103/ijcm.ijcm_616_22
11. Hadi H, Nurunniyah S, Gittelsohn J, et al. Preconception Maternal Mentoring for Improved Fetal Growth among Indonesian Women: Results from a Cluster Randomized Controlled Trial. *Nutrients.* 2023;15(21). doi:10.3390/nu15214579
12. Sari K, Sartika RAD. The Effect of the Physical Factors of Parents and Children on Stunting at Birth Among Newborns in Indonesia. *J Prev Med Public Health.* 2021;54(5):309-16. doi:10.3961/jpmph.21.120

13. Siswati T, Hookstra T, Kusnanto H. Stunting among children Indonesian urban areas: What is the risk factors? *Jurnal Gizi dan Dietetik Indonesia (Indones J Nutr Diet)*. 2020;8(1):1. doi:10.21927/ijnd.2020.8(1).1-8
14. Faienza MF, D'Amato E, Natale MP, et al. Metabolic Bone Disease of Prematurity: Diagnosis and Management. *Front Pediatr*. 2019;7. doi:10.3389/fped.2019.00143
15. Cuestas E, Hillman M, Galetto S, et al. Inflammation induces stunting by lowering bone mass via GH/IGF-1 inhibition in very preterm infants. *Pediatr Res*. 2023;94(3):1136-44. doi:10.1038/s41390-023-02559-5
16. Arroyo Pino V, Bancalari Molina A. Low levels of insulin-like growth factor 1 (IGF-1) and morbidity in preterm newborns. *Andes Pediatrica*. 2023;94(4):453-461. doi:10.32641/andespaeiatr.v94i4.4514
17. Smerieri A, Petraroli M, Ziveri MA, Volta C, Bernasconi S, Street ME. Effects of cord serum insulin, IGF-II, IGFBP-2, IL-6 and cortisol concentrations on human birth weight and length: pilot study. *PLoS One*. 2011;6(12):e29562. doi:10.1371/journal.pone.0029562
18. Pankiewicz K, Szczerba E, Maciejewski T, Fijałkowska A. Non-obstetric complications in preeclampsia. *Przeglad Menopausalny*. Termedia Publishing House Ltd. 2019;18(2):99-109. doi:10.5114/pm.2019.85785
19. Tanner MS, Malhotra A, Davey MA, Wallace EM, Mol BW, Palmer KR. Maternal and neonatal complications in women with medical comorbidities and preeclampsia. *Preg Hypertens*. 2022;27:62-68. doi:10.1016/j.prehy.2021.12.006
20. Pramana C, Peranawengrum KB, Juliani V, et al. Maternal characteristics and perinatal outcomes in women with severe preeclampsia. *Systematic Reviews in Pharmacy*. 2020;11(11):549-53. doi:10.31838/srp.2020.11.80
21. Hosny AEDMS, Fakhry MN, El-Khayat W, Kashef MT. Risk factors associated with preterm labor, with special emphasis on preterm premature rupture of membranes and severe preterm labor. *J Chinese Med Assoc*. 2020;83(3):280-7. doi:10.1097/JCMA.0000000000000243
22. Nurfaizah A, Silvana R, Dwirianti R. Association between Urinary Tract Infection and Premature Rupture of Membrane in Muhammadiyah Palembang Hospital. 2020;1. doi:10.32502/msj.v1i1.2612
23. Lu Q, Zhang X, Wang Y, et al. Sleep disturbances during pregnancy and adverse maternal and fetal outcomes: A systematic review and meta-analysis. *Sleep Med Rev*. W.B. Saunders Ltd. 2021;58. doi:10.1016/j.smrv.2021.101436
24. Muralikrishnan Nambiar, Nikhil shetty, Athulya sreenivas, Anupama suresh Y, Anjali Suneel. Human Immunodeficiency Virus in Pregnancy a Retrospective Study on Maternal and Perinatal Outcomes. *Indones J Obstet Gynecol*. Published online May 20, 2024;72-8. doi:10.32771/inajog.v12i2.2016
25. Veerappan PR, Lenin D, Sahaya Sona Thresa. Prevalence of Gestational Diabetes and its Related Risk Factors among Rural Pregnant Women. *Indones J Obstet Gynecol*. Published online May 20, 2024;79-84. doi:10.32771/inajog.v12i2.1822
26. Preda A, Pădureanu V, Moța M, et al. Analysis of maternal and neonatal complications in a group of patients with gestational diabetes mellitus. *Medicina (Lithuania)*. 2021;57(11). doi:10.3390/medicina57111170
27. Ahinkorah BO. Maternal age at first childbirth and under-five morbidity in sub-Saharan Africa: analysis of cross-sectional data of 32 countries. *Archives of Public Health*. 2021;79(1). doi:10.1186/s13690-021-00674-5
28. Londero AP, Rossetti E, Pittini C, Cagnacci A, Driul L. Maternal age and the risk of adverse pregnancy outcomes: A retrospective cohort study. *BMC Pregnancy Childbirth*. 2019;19(1). doi:10.1186/s12884-019-2400-x
29. Amaha ND, Woldeamanuel BT. Maternal factors associated with moderate and severe stunting in Ethiopian children: analysis of some environmental factors based on 2016 demographic health survey. *Nutr J*. 2021;20(1):18. doi:10.1186/s12937-021-00677-6
30. Gonete AT, Kassahun B, Mekonnen EG, Takele WW. Stunting at birth and associated factors among newborns delivered at the University of Gondar Comprehensive Specialized Referral Hospital. *PLoS One*. 2021;16(1):e0245528. doi:10.1371/journal.pone.0245528
31. Girma S, Fikadu T, Agdew E, et al. Factors associated with low birthweight among newborns delivered at public health facilities of Nekemte town, West Ethiopia: a case control study. *BMC Pregnancy Childbirth*. 2019;19(1):220. doi:10.1186/s12884-019-2372-x
32. Cai C, Vandermeer B, Khurana R, et al. The impact of occupational activities during pregnancy on pregnancy outcomes: a systematic review and metaanalysis. *Am J Obstet Gynecol*. Mosby Inc. 2020;222(3):224-38. doi:10.1016/j.ajog.2019.08.059
33. Cai C, Vandermeer B, Khurana R, et al. The impact of occupational shift work and working hours during pregnancy on health outcomes: a systematic review and meta-analysis. *Am J Obstet Gynecol* Mosby Inc. 2019;221(6):563-76. doi:10.1016/j.ajog.2019.06.051
34. Mozurkewich E. Working conditions and pregnancy outcomes: an updated appraisal of the evidence. *Am J Obstet Gynecol*. 2020;222(3):201-3. doi:10.1016/j.ajog.2019.11.1263
35. Mousa A, Naqash A, Lim S. Macronutrient and Micronutrient Intake during Pregnancy: An Overview of Recent Evidence. *Nutrients*. 2019;11(2):443. doi:10.3390/nu11020443
36. Diana S, Wahyuni CU, Prasetyo B. Maternal Complications and Risk Factors for Mortality. 2020; 9; 2020. doi:10.4081/jphr.2020.1842
37. Young MF, Nguyen PH, Gonzalez Casanova I, et al. Role of maternal preconception nutrition on offspring growth and risk of stunting across the first 1000 days in Vietnam: A prospective cohort study. *PLoS One*. 2018;13(8):e0203201. doi:10.1371/journal.pone.0203201
38. Thahir AIA, Li M, Holmes A, Gordon A. Exploring Factors Associated with Stunting in 6-Month-Old Children: A Population-Based Cohort Study in Sulawesi, Indonesia. *Nutrients*. 2023;15(15). doi:10.3390/nu15153420
39. Muglia LJ, Benhalima K, Tong S, Ozanne S. Maternal factors during pregnancy influencing maternal, fetal, and childhood outcomes. *BMC Med*. BioMed Central Ltd. 2022;20(1). doi:10.1186/s12916-022-02632-6
40. Toma TM, Andargie KT, Alula RA, Kebede BM, Gujo MM. Factors associated with wasting and stunting among children aged 06–59 months in South Ari District, Southern Ethiopia: a community-based cross-sectional study. *BMC Nutr*. 2023;9(1). doi:10.1186/s40795-023-00683-3

41. Wand H, Naidoo S, Govender V, Reddy T, Moodley J. Preventing Stunting in South African Children Under 5: Evaluating the Combined Impacts of Maternal Characteristics and Low Socioeconomic Conditions. *J Prev.* 2024;45(3):339-55. doi:10.1007/s10935-024-00766-2
42. Arabzadeh H, Doosti-Irani A, Kamkari S, Farhadian M, Elyasi E, Mohammadi Y. The maternal factors associated with infant low birth weight: an umbrella review. *BMC Pregnancy Childbirth.* 2024;24(1). doi:10.1186/s12884-024-06487-y
43. Pereira PP da S, Da Mata FAF, Figueiredo ACG, de Andrade KRC, Pereira MG. Maternal Active Smoking During Pregnancy and Low Birth Weight in the Americas: A Systematic Review and Meta-analysis. *Nicotine & Tobacco Research.* 2017;19(5):497-505. doi:10.1093/ntr/ntw228
44. Nair M, S. G, Yakoob R, N. C. C. Effect of maternal anaemia on birth weight of term babies. *Int J Contemp Pediatrics.* 2018;5(3):1019. doi:10.18203/2349-3291. ijcp20181533
45. Shifti DM, Chojenta C, Holliday EG, Loxton D. Maternal anemia and baby birth size mediate the association between short birth interval and under-five undernutrition in Ethiopia: a generalized structural equation modeling approach. *BMC Pediatr.* 2022;22(1). doi:10.1186/s12887-022-03169-6
46. Bukhari IA, Alzahrani NM, Alanazi GA, Al-Taleb MA, AlOtaibi HS. Anemia in Pregnancy: Effects on Maternal and Neonatal Outcomes at a University Hospital in Riyadh. *Cureus.* Published online July 25, 2022. doi:10.7759/cureus.27238
47. Padonou SGR, Aguemon B, Bognon GMA, et al. Poor maternal anthropometric characteristics and newborns' birth weight and length: A cross-sectional study in Benin. *Int Health.* 2019;11(1):71-77. doi:10.1093/inthealth/ihy056
48. Black RE, Victora CG, Walker SP, et al. Maternal and child undernutrition and overweight in low-income and middle-income countries. *The Lancet.* Elsevier B.V. 2013;382(9890):427-451. doi:10.1016/S0140-6736(13)60937-X